

SUPPLEMENTAL DATA**Efficacy and safety of CalliSpheres drug-eluting bead
bronchial arterial infusion chemoembolization vs. bland
embolization in advanced lung cancer with hemoptysis: A
multicenter retrospective study**

**Song Liu^{1#}, Maoli Yin^{1#}, Song Liu², Huichao Xu³, Guangji Yu², Xianchuang Liu¹,
Guimin Chen^{2*}, Weiwei Zhang^{1*}**

¹Department of Interventional Radiology, Affiliated Hospital of Jining Medical College, Jining, China;

²Tumor Center, Linyi Tumor Hospital, Linyi, China;

³Endoscopic Minimally Invasive Diagnosis and Treatment Center, Linyi People's Hospital Intersection of Wohushan Road and Wuhan Road in Beicheng New Area, Linyi, China.

*Correspondence to Weiwei Zhang: 18678766717@163.com and Guimin Chen: chengguiminlyszlyy@163.com.

[#]Contributed equally to this work: Song Liu and Maoli Yin.

Full article is available at the following link: [Efficacy and safety of CalliSpheres drug-eluting bead bronchial arterial infusion chemoembolization vs. bland embolization in advanced lung cancer with hemoptysis: A multicenter retrospective study](#)

Supplementary table 1. Data on the hemoptysis-free survival curves of the two groups of patients

First-order control		Interval start time	Number of access intervals	Number of cancellations during the time interval	Number at risk	Number of terminal events	Percentage of termination	Survival analysis ratio	Proportion of cumulative survival analysis systems at the end of the period	Standard error of the cumulative survival analysis ratio at the end of the period	Probability density	Standard error of probability density	Risk rate	Standard error of risk ratio
Groups	Observation Group	0	76	0	76	0	0.00	1.00	1.00	0.00	0.000	0.000	0.00	0.00
		25	76	0	76	0	0.00	1.00	1.00	0.00	0.000	0.000	0.00	0.00
		50	76	0	76	10	0.13	0.87	0.87	0.04	0.005	0.002	0.01	0.00
		75	66	0	66	38	0.58	0.42	0.37	0.06	0.020	0.002	0.00	0.00

													3	
		100	28	0	28	22	0.79	0.21	0.08	0.03	0.012	0.002	0.05	0.01
		125	6	0	6	6	1.00	0.00	0.00	0.00	0.003	0.001	0.08	0.00
	Control subjects	0	68	0	68	0	0.00	1.00	1.00	0.00	0.000	0.000	0.00	0.00
		25	68	0	68	32	0.47	0.53	0.53	0.06	0.019	0.002	0.02	0.00
		50	36	0	36	30	0.83	0.17	0.09	0.03	0.018	0.002	0.06	0.01
		75	6	0	6	6	1.00	0.00	0.00	0.00	0.004	0.001	0.08	0.00

Supplementary table 2. Data on the overall survival curves of the two groups of patients

First-order control		Interval start time	Number of access intervals	Number of cancellations during the time interval	Number at risk	Number of terminal events	Percentage of termination	Survival analysis ratio	Proportion of cumulative survival analysis systems at the end of the period	Standard error of the cumulative survival analysis ratio at the end of the period	Probability density	Standard error of probability density	Risk rate	Standard error of risk ratio
Groups	Observation Group	0	76	0	76	7	0.09	0.91	0.91	0.03	0.018	0.007	0.02	0.01
		5	69	0	69	29	0.42	0.58	0.53	0.06	0.076	0.011	0.11	0.02
		10	40	0	40	16	0.40	0.6	0.32	0.05	0.042	0.009	0.10	0.03
		15	24	0	24	15	0.63	0.38	0.12	0.04	0.039	0.009	0.1	0.04

													8	
		20	9	2	8	4	0.50	0.5	0.06	0.03	0.012	0.006	0.13	0.06
		25	3	3	2	0	0.00	1	0.06	0.03	0.000	0.000	0.00	0.00
	Control subjects	0	68	0	68	12	0.18	0.82	0.82	0.05	0.035	0.009	0.04	0.01
		5	56	0	56	45	0.80	0.2	0.16	0.04	0.132	0.011	0.27	0.03
		10	11	0	11	11	1.00	0	0.00	0.00	0.032	0.009	0.40	0.00