

# Can NLR, PLR, and LMR be used as prognostic indicators in patients with pulmonary embolism? A commentary

Cihan Bedel<sup>1\*</sup>, Mustafa Korkut<sup>1</sup>, Hamit Hakan Armağan<sup>2</sup>

We read with great interest the article “Prognostic role of neutrophil-lymphocyte ratio (NLR), platelet-lymphocyte ratio (PLR), and lymphocyte-monocyte ratio (LMR) in patients with pulmonary embolism” by Köse et al. [1]. They found that the NLR, PLR, and LMR were related to the prognosis and clinical severity of patients with pulmonary embolism (PE). First of all, we congratulate the authors for their invaluable contribution to literature. However, we think that there are some points that should be discussed regarding the use of these laboratory parameters.

White blood cell subtypes NLR, PLR, and LMR have been associated with many inflammatory diseases, including PE [2,3]. These parameters, which can be easily determined by simple and easy measurement of systemic inflammation, maintain their importance today. However, these parameters are affected by many factors such as trauma, local or systemic infection, acute coronary syndromes, and malignancy [3-5]. For these reasons, it would be better for the authors to mention these factors and exclude them from the tables that included malignancy and trauma patients in the study.

It is known that drugs, including steroids, can increase neutrophils and decrease lymphocytes and therefore affect NLR, PLR, and LMR values [6]. Therefore, it will be more valuable to exclude patients who use drugs that may affect laboratory parameters. In addition, plasma inflammatory biomarkers are time-dependent variables, the time of sample collection and the time from the onset of the symptom to the sampling may have an impact on the parameters [3-6]. Therefore, it is

important to identify the time from the first symptom to sample collection and the factors that may affect it. In conclusion, because NLR, PLR, and LMR can be affected by many factors, prospective studies with large populations are needed to show the accuracy of use in critically ill patients.

## REFERENCES

1. Köse N, Yıldırım T, Akın F, Yıldırım SE, Altun İ. Prognostic role of NLR, PLR, and LMR in patients with pulmonary embolism. *Bosn J Basic Med Sci* 2020;20(2):248-53. <https://doi.org/10.17305/bjbms.2019.4445>.
2. Bedel C, Selvi F. Association of platelet to lymphocyte and neutrophil to lymphocyte ratios with in-hospital mortality in patients with Type A acute aortic dissection. *Braz J Cardiovasc Surg* 2019;34(6):694-8. <https://doi.org/10.21470/1678-9741-2018-0343>.
3. Cho KH, Jeong MH, Ahmed K, Hachinohe D, Choi HS, Chang SY, et al. Value of early risk stratification using hemoglobin level and neutrophil-to-lymphocyte ratio in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. *Am J Cardiol* 2011;107(6):849-56. <https://doi.org/10.1016/j.amjcard.2010.10.067>.
4. Huszno J, Kolosza Z. Prognostic value of the neutrophil-lymphocyte, platelet-lymphocyte and monocyte-lymphocyte ratio in breast cancer patients. *Oncol Lett* 2019;18(6):6275-83. <https://doi.org/10.3892/ol.2019.10966>.
5. Özkaya M, Bedel C. Can neutrophil to lymphocyte ratio and platelet to lymphocyte ratio predict the severity of sternum fractures? *Curr Thorac Surg* 2018;3(3):116-22. <https://doi.org/10.26663/cts.2019.00022>.
6. Bedel C, Korkut M. Comment on: Neutrophil to lymphocyte ratio is associated with in-hospital mortality in older adults admitted to the emergency department. *Am J Emerg Med* 2020;In press. <https://doi.org/10.1016/j.ajem.2020.10.001>.

<sup>1</sup>Department of Emergency Medicine, Health Science University Antalya Training and Research Hospital, Antalya, Turkey

<sup>2</sup>Department of Emergency Medicine, Faculty of Medicine, Suleyman Demirel University, Isparta, Turkey

\*Corresponding author: Cihan Bedel, Department of Emergency Medicine, Health Science University Antalya Training and Research Hospital, Kazım Karabekir Street, Postal Zip Code: 07100, Muratpaşa, Antalya, Turkey. Phone: +905075641254, Fax: +902422494487. E-mail: cihanbedel@hotmail.com

DOI: <https://doi.org/10.17305/bjbms.2020.5219> 20.5236

Submitted: 14 October 2020/ Accepted: 14 October 2020

Conflicts of interest statement: The authors declare no conflicts of interest.



©The Author(s) (2020). This work is licensed under a Creative Commons Attribution 4.0 International License

## Related articles published in BJBMS

1. [Prognostic role of NLR, PLR, and LMR in patients with pulmonary embolism](#)  
Nuri Köse et al., BJBMS, 2019
2. [Can NLR, PLR and LMR be used as prognostic indicators in patients with pulmonary embolism? Author's reply on commentary](#)  
Nuri Kose et al., BJBMS, 2020
3. [Prognostic role of neutrophil-to-lymphocyte ratio \(NLR\) in patients with operable ampullary carcinoma](#)  
Nebi Serkan Demirci et al., BJBMS, 2018